

Resonance neutrino bremsstrahlung $\nu \rightarrow \nu\gamma$ in a strong magnetic field

A.A. Gvozdev, N.V. Mikheev

*Division of Theoretical Physics, Department of Physics,
Yaroslavl State University, Yaroslavl 150000, Russia
E-mail: mikheev@yars.free.net*

and

L.A. Vassilevskaya

*Moscow Lomonosov University, V-952, Moscow 117234, Russia
E-mail: vasilevs@vitep5.itep.ru*

Abstract

High energy neutrino bremsstrahlung $\nu \rightarrow \nu + \gamma$ in a strong magnetic field ($B \gg B_s$) is studied in the framework of the Standard Model (SM). A resonance probability and a four-vector of the neutrino energy and momentum loss are presented. A manifestation of the neutrino bremsstrahlung in astrophysical cataclysm of type of a supernova explosion or a merger of neutron stars, such as an possible origin of cosmological γ -burst is discussed.

Neutrino physics in media is a vigorously growing and prospective line of investigation at the junction of particle physics, astrophysics and cosmology. Substance is usually considered as the medium. We stress that a strong magnetic field can play role of the active medium. It will suffice to mention that the magnetic field can influence substantially on neutrino electromagnetic properties (an anomalous magnetic moment [1], neutrino oscillations [2], a magnetic catalysis of the radiative decay [3]). The investigations of this sort are, in our opinion, of principal interest in astrophysics where gigantic neutrino fluxes and strong magnetic fields ($B > B_s$, $B_s = m_e^2/e \simeq 0.44 \cdot 10^{14} G$ is the critical, so called, Schwinger value) can take place simultaneously (a process of a coalescence of neutron stars, an explosion of a supernova). Notice that the magnetic fields at the moment of cataclysm can occur much more than the typical, so called, "old" magnetic fields of order of B_s which have been observed at the surface of pulsars. By now astrophysical knowledge of strong magnetic fields which can be realized in the nature have changed essentially. For example, the field strengths inside the astrophysical objects in principle could be as high as $10^{15} - 10^{18} G$, both for toroidal [4] and for poloidal [5] fields. On the other hand, in such strong magnetic fields otherwise negligible processes are not only opened kinematically but become substantial ones as well.

In our paper [3] we have studied the massive neutrino radiative decay $\nu \rightarrow \nu' + \gamma$ [6] in external electromagnetic fields of various configurations, in particular, in the strong magnetic field. It was shown that a field-induced amplitude was not suppressed by smallness of neutrino masses and did not vanish even in the case of massless neutrino as opposed to the vacuum amplitude. The decay probability of the neutrino with energy $E_\nu < 2m_e$ was calculated on the assumption that photon dispersion relation was close to the vacuum one $q^2 = 0$.

However the photon dispersion in the strong magnetic field differs significantly from the vacuum dispersion with increasing photon energy [7–9], so the real photon 4-momentum can appear as a space-like and sufficiently large one ($|q^2| \gg m_\nu^2$). In this case the phase space for neutrino transition $\nu \rightarrow \nu' + \gamma$ with $m < m'$ is opened also. It means that the decay probability of ultrarelativistic neutrino $\nu \rightarrow \nu' + \gamma$ becomes insensitive to the neutrino mass spectrum due to the photon dispersion relation in the strong magnetic field. This phenomenon results in a strong suppression ($\sim m_\nu^2/E_\nu^2$) of the neutrino transition with flavour violation and a diagonal process $\nu_l \rightarrow \nu_l + \gamma$ ($l = e, \mu, \tau$) is realized only. Thus this diagonal radiative neutrino transition manifests itself as neutrino bremsstrahlung, does not contain uncertainties associated with a possible mixing in the lepton sector of the SM and can lead to observable physical effects in the strong magnetic fields $B \gg B_s$.

In this paper we study the high energy neutrino bremsstrahlung in the strong constant magnetic field in the framework of the SM. The field-induced $\nu\nu\gamma$ -vertex

can be calculated using an effective four-fermion weak interaction of the left neutrino with the electron only because the electron is the most sensitive fermion to the external field. By this means diagrams describing this process are reduced to an effective diagram with the electron in the loop (Fig.1)




FIG. 1.

where double lines imply that the influence of the external field in propagators is taken into account exactly.

The calculation technique of this type loop diagram was described in detail in our paper [3]. The general expression for the $\nu\nu\gamma$ -vertex has a cumbersome form and will be published elsewhere. We note that it is enhanced substantially in the vicinity of the, so called, photon cyclotron frequencies similar to a vacuum polarization (the cyclotron resonance in the vacuum polarization [10]). Here we present the amplitude of the high energy neutrino ($E_\nu \gg m_e$) bremsstrahlung $\nu_l(p) \rightarrow \nu_l(p') + \gamma(q)$ in the case when the field strength is the greatest parameter $eB > E_\nu^2$. We stress that two eigenmodes of the photon propagation with polarization vectors

$$\varepsilon_\mu^{(\parallel)} = \frac{(q\varphi)_\mu}{\sqrt{q_\perp^2}}; \quad \varepsilon_\mu^{(\perp)} = \frac{(q\tilde{\varphi})_\mu}{\sqrt{q_\parallel^2}} \quad (1)$$

are realized in the magnetic field (the, so called, parallel ($\parallel$) and perpendicular ($\perp$) polarizations according to Adler's notations in the paper [7]). Here $\varphi_{\alpha\beta} = F_{\alpha\beta}/B$ and $\tilde{\varphi}_{\alpha\beta} = \frac{1}{2}\varepsilon_{\alpha\beta\mu\nu}\varphi_{\mu\nu}$ are the dimensionless tensor and dual tensor of the external magnetic field with the strength $\vec{B} = (0, 0, B)$; $q_\parallel^2 = (q\tilde{\varphi}\tilde{\varphi}q) = q_\alpha\tilde{\varphi}_{\alpha\beta}\tilde{\varphi}_{\beta\mu}q_\mu = q_0^2 - q_3^2$, $q_\perp^2 = (q\varphi\varphi q) = q_1^2 + q_2^2$. With (1) the amplitude $M = M_\parallel + M_\perp$ is significantly simplified and can be presented in the form:

$$M_\parallel \simeq -\frac{eG_F}{\sqrt{2}} g_v V_\parallel \frac{(q\varphi j)}{\sqrt{q_\perp^2}} \quad (2)$$

$$M_\perp \simeq -\frac{eG_F}{\sqrt{2}} \frac{1}{\sqrt{q_\parallel^2}} \{g_v V_\perp (q\tilde{\varphi} j) + g_a A(q\varphi\varphi j)\},$$

where $M_\parallel$ and $M_\perp$ correspond to the creation of the $\parallel$ and $\perp$ photon modes, respectively. Here $e > 0$ is the elementary charge; G_F is the Fermi constant, g_v and g_a are the vector and axial-vector $\nu - e$ coupling constants

in the SM, respectively; $j_\alpha = 2\bar{\nu}_L(p')\gamma_\alpha\nu_L(p)$ is the left neutrino current, $V_\parallel$, $V_\perp$ and A are the scalar functions of $q_\parallel^2$, $q_\perp^2$ and eB . Notice that $V_\parallel$ and $V_\perp$ are the known functions (see, for example, [10]) since the field-induced vacuum polarization is described in terms of these functions, namely:

$$\mathcal{P}_\lambda = e^2\pi(q^2) + e^2 V_\lambda, \quad \lambda = \parallel, \perp, \quad (3)$$

where $\pi(q^2)$ is the well-known vacuum polarization without an external field ($q^2 = q_\parallel^2 - q_\perp^2$). As for the function A the result of our calculations is:

$$\begin{aligned} A &= \frac{1}{4\pi^2} \frac{q_\parallel^2 + q_\perp^2}{q_\perp^2} \left[i \int_0^1 du \int_0^\infty dt e^{-i\Phi} \right. \\ &\quad \times \left(m_e^2 - \frac{q_\parallel^2 \cdot q^2}{q_\parallel^2 + q_\perp^2} \frac{1-u^2}{4} \right) - eB \Big], \\ \Phi &= \frac{1}{eB} \left[t \left(m_e^2 - q_\parallel^2 \frac{1-u^2}{4} \right) + q_\perp^2 \frac{\cos ut - \cos t}{2 \sin t} \right], \end{aligned} \quad (4)$$

The important properties of functions V_λ and A are that they are singular with respect to the variable $q_\parallel^2$ in the threshold points $(q_\parallel^2)_{thr} = \mathcal{E}_{nm}^2$, labelled by n, m :

$$\mathcal{E}_{nm}^2 = \left(\sqrt{m_e^2 + 2eBn} + \sqrt{m_e^2 + 2eBm} \right)^2, \quad (5)$$

where $\mathcal{E}_{nm}$ has the simple physical meaning as transversal energy of an electron-positron pair in the magnetic field.

It is of interest for some astrophysical applications the case of relatively high neutrino energy $E_\nu \simeq 10 - 20 MeV \gg m_e$ and strong magnetic field $eB > E_\nu^2$ when a region of the cyclotron resonance on the ground Landau level dominates in this process. The functions $V_\perp$ [10] and A are easily calculated in this region:

$$A \simeq -V_\perp \simeq \frac{eB m_e e^{-q_\perp^2/2eB}}{2\pi\sqrt{4m_e^2 - q_\parallel^2}}. \quad (6)$$

As it is seen from (6) the amplitude $M_\perp$ (2) contains the enhancement due to the square-root singularity which is connected with the cyclotron resonance on the ground Landau level ($q_\parallel^2 \rightarrow 4m_e^2$). As for $V_\parallel$, it has not such an enhancement and the dispersion relation for $\parallel$ photon mode is close to the vacuum one. So, the amplitude $M_\parallel \sim (q\varphi j) \sim q^2$ is small due to collinear kinematics of the process with creation of $\parallel$ photon mode. The square-root singularities on higher Landau levels are not realized under the condition $eB > E_\nu^2$. We note that not only function A has singular behaviour in the vicinity of the threshold point $q_\parallel^2 = 4m_e^2$, but the vacuum polarization $\mathcal{P}_\perp$ (3) as well. This means that many-loops quantum corrections are of great importance in the vicinity of the resonance. As analysis shows taking account of these radiative corrections reduces to the renormalization of the $\perp$ photon mode wave function:

$$\varepsilon_\mu^{(\perp)} \rightarrow \sqrt{Z} \varepsilon_\mu^{(\perp)}, \quad Z = \left(1 - \frac{\partial \mathcal{P}_\perp}{\partial q_\parallel^2} \right)^{-1}, \quad (7)$$

and leads to an additional factor $\sqrt{Z}$ in the amplitude $M_\perp$ (2).

The probability of the neutrino bremsstrahlung $\nu \rightarrow \nu\gamma$ can be obtained by integration over the phase space with taking account of the photon $q^2 - \mathcal{P}_\perp = 0$ in the vicinity of the resonance:

$$4m_e^2 - q_\parallel^2 \simeq \alpha^2 \left(\frac{2eB}{q_\perp^2} \right)^2 m_e^2 \ll q_\perp^2.$$

This leads to the dependence of photon energy on the photon momentum $q_0 \simeq \sqrt{q_3^2 + 4m_e^2}$ (the photon created can move only along the magnetic field with the velocity $v = dq_0/dq_3 \simeq q_3/\sqrt{q_3^2 + 4m_e^2}$). The main contribution to the probability is determined from the vicinity of the resonance point $q_\parallel^2 = 4m_e^2$ and can be presented in the form:

$$\begin{aligned} W &\simeq \frac{\alpha G_F^2}{8\pi^2} (g_v^2 + g_a^2) (eB)^2 E_\nu \sin^2 \theta F(\lambda), \\ F(\lambda) &= 1 - \frac{\lambda}{2} + \frac{\lambda^2}{3} - \frac{5\lambda^3}{24} + \frac{7\lambda^4}{60} + O(\lambda^5), \\ \lambda &= \frac{E_\nu^2 \sin^2 \theta}{eB} < 1, \end{aligned} \quad (8)$$

where θ is the angle between the vectors of the magnetic field strength $\vec{B}$ and the momentum of the initial neutrino $\vec{p}$.

We stress the principal importance of taking account of the photon wave function renormalization (7). If one ignored this factor the decay probability would be strongly overstated:

$$W \simeq \left(\frac{1}{\alpha} \right) \left(\frac{5G_F^2 m_e^5}{256\pi^2} \right) (g_v^2 + g_a^2) \left(\frac{E_\nu}{m_e} \right)^7 \sin^8 \theta. \quad (9)$$

It is of more practical importance for some astrophysical applications to calculate the four-vector of the neutrino energy and momentum loss in the process of the neutrino bremsstrahlung:

$$Q_\mu = E_\nu \int dW \cdot q_\mu. \quad (10)$$

The physical meaning of this vector is that Q_0/E_ν is the mean energy loss in a unit time and $\vec{Q}/E_\nu$ is the momentum loss in a unit time. We have obtained the expression for the vector Q_μ :

$$\begin{aligned} Q_\mu &= \frac{1}{4} E_\nu W \left(F_1(\lambda)(p\tilde{\Lambda})_\mu - 2F_2(\lambda)(p\Lambda)_\mu \right. \\ &\quad \left. - \frac{2g_v g_a}{g_v^2 + g_a^2} F_1(\lambda)(p\tilde{\varphi})_\mu \right), \\ F_1(\lambda) &= 1 - \frac{2\lambda}{3} + \frac{5\lambda^2}{12} - \frac{7\lambda^3}{30} + \frac{7\lambda^4}{60} + O(\lambda^5), \\ F_2(\lambda) &= 1 - \lambda + \frac{5\lambda^2}{6} - \frac{7\lambda^3}{12} + \frac{7\lambda^4}{20} + O(\lambda^5). \end{aligned} \quad (11)$$

To illustrate the applications of the results we have obtained we estimate below possible effects of the neutrino bremsstrahlung in astrophysical cataclysm of type of a supernova explosion or a merger of neutron stars under the condition that a compact remnant has for some reasons extremely strong magnetic field, $B \simeq 10^{16} - 10^{18} G$. We assume that neutrinos of all species with the typical mean energy $\bar{E}_\nu \simeq 20 MeV$ are radiated from a surface of a neutrinosphere in such a process. Using (11) we estimate the neutrino energy loss:

$$\mathcal{E} \simeq 10^{50} \left(\frac{\mathcal{E}_{tot}}{10^{53} erg} \right) \left(\frac{B}{10^{17} G} \right)^2 \times \left(\frac{\bar{E}_\nu}{20 MeV} \right) \left(\frac{R}{10 km} \right) erg, \quad (12)$$

where $\mathcal{E}_{tot} \simeq 10^{53}$ erg is the total neutrino radiation energy [11]; B is the magnetic field strength in the vicinity of the neutrinosphere of radius R . Neutrino bremsstrahlung γ -quanta are captured by a strong magnetic field and propagate along the field [10]. Thus the mechanism of significant power "pumping" of polar caps of a magnetized remnant can take place. This phenomenon can result in reemission within a narrow cone $\Omega/4\pi \ll 1$ along the magnetic moment of the remnant. So, if an external absorbing envelope is absent in such an extraordinary astrophysical cataclysm (very strong magnetic fields, gigantic neutrino fluxes) the reemission process will be an observable effect. Namely, such a reemission in the magnetic field could appear as an anisotropic γ -burst with the duration of order of the neutrino emission time and of the typical energy $\mathcal{E}_{4\pi}^{burst} \sim 10^{51}$ erg in terms of 4π -geometry.

The authors thank V.A. Rubakov and V.M. Lipunov for fruitful discussions, M.E. Shaposhnikov and A.V. Kuznetsov for useful critical remarks. The work of N.V. Mikheev was supported by a Grant N d104 from the International Soros Science Education Program. The work of L.A. Vassilevskaya was supported by a fellowship of INTAS Grant 93-2492-ext and was carried out within the research program of International Center for Fundamental Physics in Moscow.

-
- [1] A.V. Borisov et al., *Yad. Fiz.* **41** 743 (1985).
 - [2] J.F. Nieves and P.B. Pal, *Phys. Rev. D* **40** 1693 (1989); J.C. D'Olivo, J.F. Nieves and P.B. Pal, *Phys. Rev. D* **40** 3679 (1989).
 - [3] A.A. Gvozdev, N.V. Mikheev and L.A. Vassilevskaya, *Phys. Rev. D* **54** 5674 (1996).
 - [4] M. Ruderman, in "Neutron Stars: Theory and Observation", ed. by J. Ventura and D. Pines, Kluwer Academic. Pub., Dordrecht, 1991; V.M. Lipunov, *Astrophysics of Neutron Stars*, Springer-Verlag, New York, 1992, and V.M. Lipunov, private communication.
 - [5] M. Bocquet et al., *Astr. and Ap.* **301** 757 (1995); V.V.

- Usov, *Nature* **357** 472 (1992); R. Narayan, B. Paczynski and T. Piran, *Ap.J.* **395** L83 (1992).
- [6] We stress that ν and ν' are the neutrinos eigenstates with definite masses, but not the neutrinos with definite flavours.
- [7] S.L. Adler, *Ann. Phys. N.Y.* **67** 599 (1971);
- [8] I.A. Batalin and A.E. Shabad, *Sov.Phys.JETP* **33** 483 (1971).
- [9] W.-Y. Tsai, *Phys. Rev. D* **10** 2699 (1974).
- [10] A.E. Shabad, *Ann. Phys. N.Y.* **90** 166 (1975); A.E. Shabad and V.V. Usov, *Astrophysics and Space Science*. **102** 327 (1984).
- [11] H.T. Janka and M. Ruffert, *Astr. and Ap.* **307** L33 (1996); V.S. Imshennik and D.K. Nadyozhin, *Sov. Sci. Rev., Sect. E* 8(1989)1; D.K. Nadyozhin, in: *Particles and Cosmology*, Proc. Baksan Int. School, eds. V.A. Matveev et al., World Sci., 1992, p.153.